

DIAGNOSIS AND MANAGEMENT OF CERVICAL INSUFFICIENCY

DEFINITION AND INTRODUCTION

Cervical insufficiency is defined as the inability of the uterine cervix to retain a pregnancy in the second trimester, in the absence of uterine contractions.⁽¹⁾

A history of cervical insufficiency has been applied to women with one or more second trimester pregnancy losses/preterm births (before 34 weeks) who fulfil this definition. It must be noted that a short cervical length on transvaginal scan in the second trimester is a risk factor for preterm birth but is not sufficient to diagnose cervical insufficiency.

Prematurity is the leading cause of perinatal death and disability. Evidence suggests that the incidence of preterm labour and birth is continuing to rise worldwide. Currently 6% of babies in New Zealand are born preterm. Despite efforts and interventions aimed at reducing the incidence globally the results have been largely disappointing.⁽²⁾

It can be difficult to distinguish between women who have a short cervix and those that have true cervical insufficiency.

RISK FACTORS

Refer to the Obstetric clinic is guided by Section 88 Referral Guideline.

CERVICAL RISK FACTORS: (SEE APPENDIX A)

- **Collagen abnormalities** — genetic disorders affecting collagen (eg., Ehlers Danlos syndrome) have been associated with an increased risk of preterm birth⁽⁴⁾
- **Uterine anomalies** — increase the risk of second trimester preterm birth, eg. Septate uterus, bicornuate uterus and even arcuate uterus⁽⁵⁾
- **Biologic variation** — although a short cervix is predictive of preterm birth, it is not diagnostic of cervical insufficiency and many women who have a congenitally short cervix deliver at term⁽⁶⁾

PAST OBSTETRIC HISTORY: (SEE APPENDIX B)

- Recurrent mid-trimester pregnancy losses
- Previous preterm pre-labour rupture of membranes at less than 32 weeks
- Prior pregnancy with a cervical length measurement of less than 25 mm prior to 27 weeks of gestation⁽³⁾

ACQUIRED FACTORS (MORE COMMON)

- **Cervical trauma** — may weaken the cervix, and contribute to cervical insufficiency.
- **Mechanical dilation** — eg. dilation and curettage [D&C], dilation and evacuation [D&E], pregnancy termination, hysteroscopy.^(7,8) In women with a short cervical length and no prior preterm birth, prior cervical mechanical dilatation is one of the most common associated risk factors.
- **Treatment of cervical intraepithelial neoplasia** — LEEP may increase the risk for late preterm birth (from 34 to < 37 weeks of gestation)⁹.
- Women may have no symptoms or can present with mild symptoms, eg. painless vaginal spotting, increased vaginal discharge, premenstrual-like cramping or backache or pelvic pressure
- Women may present with these symptoms from as early as 14 to 20 weeks of gestation.

DIAGNOSIS

This is either based on history alone or in combination with transvaginal ultrasound (TVU) measurement of cervical length.

Important note:

The diagnosis of cervical insufficiency is usually limited to singleton gestations because the pathogenesis of delivery at 14 to 28 weeks in multiple gestations is usually unrelated to a weakened cervix.

MANAGEMENT

The management of these women can be divided into two main groups:

- (1) Women for whom a conservative path will be pursued
- (2) Women where it is clear that surgical intervention in the form of a cerclage is indicated.
This may be either prophylactic or therapeutic.

APPROACH TO MANAGEMENT

See pathways below.


REFERENCES

1. ACOG Practice Bulletin No.142: Cerclage for the management of cervical insufficiency. American College of Obstetricians and Gynaecologists *Obstet Gynecol.* 2014; 123 (2 Pt 1):372.
2. Green Top Guideline No. 60: Cervical Cerclage May 2011. Royal College of Obstetricians and Gynaecologists
3. Drakeley AJ, Roberts D, Alfirevic Z. Cervical stitch (cerclage) for preventing pregnancy loss in women. *Cochrane Database Syst Rev.* 2003:CD003253
4. Leduc L, Wasserstrum N. Successful treatment with the Smith-Hodge pessary of cervical incompetence due to defective connective tissue in Ehlers-Danlos syndrome. *Am J Perinatol* 1992; 9:25.
5. Chan YY, Jayaprakasan K, Tan A, et al. Reproductive outcomes in women with congenital uterine anomalies: a systematic review. *Ultrasound Obstet Gynecol* 2011; 38:371.
6. Vincenzo Berghella, MD. Cervical insufficiency Up to date May 2014
7. Johnstone FD, Beard RJ, Boyd IE, McCarthy TG. Cervical diameter after suction termination of pregnancy. *Br Med J* 1976; 1:68.
8. Romero, R, Lockwood, CJ. Pathogenesis of spontaneous preterm labor. *Creasy & Resnik's Maternal Fetal Medicine*, Creasy, RK, Resnik, R, Iams, JD, Lockwood, CJ, Moore, TR (Eds), Saunders, 2009
9. Cervical intraepithelial neoplasia: Reproductive effects of treatment. Jakobsson M, Norwitz E R. Up to date May 2014
10. Final report of the Medical Research Council/Royal College of Obstetricians and Gynaecologists multicentre randomised trial of cervical cerclage. MRC/RCOG Working Party on Cervical Cerclage. *Br J Obstet Gynaecol.* 1993; 100(6):516.
11. Prophylactic administration of progesterone by vaginal suppository to reduce the incidence of spontaneous preterm birth in women at increased risk: a randomized placebo-controlled double-blind study. da Fonseca EB, Bittar RE, Carvalho MH, Zugaib M *Am J Obstet Gynecol.* 2003; 188(2):419.
12. Prevention of recurrent preterm delivery by 17 alpha-hydroxyprogesterone caproate. Meis PJ, Klebanoff M, Thom E, Dombrowski MP, Sibai B, Moawad AH, Spong CY, Hauth JC, Miodovnik M, Varner MW, Leveno KJ, Caritis SN, Iams JD, Wapner RJ, Conway D, O'Sullivan MJ, Carpenter M, Mercer B, Ramin SM, Thorp JM, Peaceman AM, Gabbe S, National Institute of Child Health and Human Development Maternal-Fetal Medicine Units Network *N Engl J Med.* 2003;348(24):2379.
13. Efficacy of progesterone support for pregnancy in women with recurrent miscarriage. A meta-analysis of controlled trials. Daya S *Br J Obstet Gynaecol.* 1989 Mar; 96(3):275-80.
14. Progesterone and preterm birth prevention: translating clinical trials data into clinical practice. Society for Maternal-Fetal Medicine Publications Committee, with assistance of Vincenzo Berghella *Am J Obstet Gynecol.* 2012;206(5):376
15. Does transvaginal sonographic measurement of cervical length before 14 weeks predict preterm delivery in high-risk pregnancies? Berghella V, Talucci M, Desai A *Ultrasound Obstet Gynecol.* 2003; 21(2):140.
16. The rate of cervical change and the phenotype of spontaneous preterm birth Iams JD, Cebrik D, Lynch C, Behrendt N, Das A *Am J Obstet Gynecol.* 2011
17. The effect of 17 α -hydroxyprogesterone caproate on preterm birth in women with an ultrasound-indicated cerclage. Rafael TJ, Mackeen AD, Berghella V *Am J Perinatol.* 2011 May; 28(5):389-94. Epub 2011 Mar 4.
18. Cervical length for prediction of preterm birth in women with multiple prior induced abortions. Visintine J, Berghella V, Henning D, Baxter J *Ultrasound Obstet Gynecol.* 2008;31(2):198.
19. Transvaginal ultrasonography of the cervix to predict preterm birth in women with uterine anomalies. Airoidi J, Berghella V, Sehdev H, Ludmir J *Obstet Gynecol.* 2005;106(3):553.
20. Prior cone biopsy: prediction of preterm birth by cervical ultrasound. Berghella V, Pereira L, Garipey A, Simonazzi *GSOAm J Obstet Gynecol.* 2004; 191(4):1393.

21. Cerclage for short cervix on ultrasonography: meta-analysis of trials using individual patient-level data. Berghella V, Odibo AO, To MS, Rust OA, Althuisius SM *Obstet Gynecol.* 2005; 106(1):181.
22. Progesterone and the risk of preterm birth among women with a short cervix. Fonseca EB, Celik E, Parra M, Singh M, Nicolaides KH, Fetal Medicine Foundation Second Trimester Screening Group *N Engl J Med.* 2007;357(5):462.
23. Vaginal progesterone reduces the rate of preterm birth in women with a sonographic short cervix: a multicenter, randomized, double-blind, placebo-controlled trial. Hassan SS, Romero R, Vidyadhari D, Fusey S, Baxter JK, Khandelwal M, Vijayaraghavan J, Trivedi Y, Soma-Pillay P, Sambarey P, Dayal A, Potapov V, O'Brien J, Astakhov V, Yuzko O, Kinzler W, Dattel B, Sehdev H, Mazheika L, Manchulenko D, Gervasi MT, Sullivan L, Conde-Agudelo A, Phillips JA, Creasy GW, PREGNANT Trial *Ultrasound Obstet Gynecol.* 2011;38(1):18.
24. Universal cervical length screening and treatment with vaginal progesterone to prevent preterm birth: a decision and economic analysis. Cahill AG, Odibo AO, Caughey AB, Stamilio DM, Hassan SS, Macones GA, Romero R *Am J Obstet Gynecol.* 2010 Jun; 202(6):548.e1-8. Epub 2010 Jan 15.
25. Universal cervical-length screening to prevent preterm birth: a cost-effectiveness analysis. Werner EF, Han CS, Pettker CM, Buhimschi CS, Copel JA, Funai EF, Thung SF *Ultrasound Obstet Gynecol.* 2011;38(1):32.
26. Multicenter randomized trial of cerclage for preterm birth prevention in high-risk women with shortened midtrimester cervical length. Owen J, Hankins G, Iams JD, Berghella V, Sheffield JS, Perez-Delboy A, Egerman RS, Wing DA, Tomlinson M, Silver R, Ramin SM, Guzman ER, Gordon M, How HY, Knudtson EJ, Szychowski JM, Cliver S, Hauth JC *Am J Obstet Gynecol.* 2009;201(4):375.e1.
27. Pregnancy outcomes in women treated with elective versus ultrasound-indicated cervical cerclage. Guzman ER, Forster JK, Vintzileos AM, Ananth CV, Walters C, Gipson K *Ultrasound Obstet Gynecol.* 1998;12(5):323.
28. Elective cerclage vs. ultrasound-indicated cerclage in high-risk pregnancies. To MS, Palaniappan V, Skentou C, Gibb D, Nicolaides KH *Ultrasound Obstet Gynecol.* 2002;19(5):475.
29. Elective cervical cerclage versus serial ultrasound surveillance of cervical length in a population at high risk for preterm delivery. Groom KM, Bennett PR, Golara M, Thalon A, Shennan AH *Eur J Obstet Gynecol Reprod Biol.* 2004; 112(2):158.
30. Etiologies and subsequent reproductive performance of 100 couples with recurrent abortion. Phung Thi Tho, Byrd JR, McDonough PG *Fertil Steril.* 1979; 32(4):389.
31. Cervical cerclage: patient selection, morbidity, and success rates. Harger JH *Clin Perinatol.* 1983; 10(2):321.
32. Cervical length screening with ultrasound-indicated cerclage compared with history-indicated cerclage for prevention of preterm birth: a meta-analysis. Berghella V, Mackeen AD *Obstet Gynecol.* 2011; 118(1):148.

APPENDIX A CONSERVATIVE MANAGEMENT

Women with cervical risk factors for cervical insufficiency but no history of previous loss.




Ref.236960

*Discuss with NICU

APPENDIX B CONSERVATIVE MANAGEMENT

Those women with a previous 2nd trimester loss or a previous preterm delivery before 34 weeks




Ref.236962

*Discuss with NICU

APPENDIX C SURGICAL MANAGEMENT

Suspected history of cervical insufficiency is:

Three or more preterm births < 34 weeks (with progressively earlier deliveries in successive pregnancies) and/or second trimester losses



Ref.236964

**No trials have evaluated the efficiency of combination therapy

*Discuss with NICU

APPENDIX D ACUTE PRESENTATION WITH SUSPECTED CERVICAL INSUFFICIENCY


History

1. Take an incidental history to rule out infection or preterm labour
2. Maternal Observations → temperature, pulse rate, blood pressure, respiratory rate
3. Examination - abdominal palpitations (fundal height, tenderness, uterine activity)
4. Vaginal assessment - speculum examination of cervical effacement and dilation
 - ◆ Exclude SROM, bleeding, abnormal vaginal discharge
5. Digital exam ONLY if evidence of advanced dilation and birth thought imminent
→ consult with senior registrar on-call

Investigations

1. MSU for culture and sensitivity
2. HVS and vulvo-vaginal swab for Chlamydia and Gonorrhoea
3. FBC, CRP
4. If visual signs of dilation and effacement consider a TVS for cervical length and TAS for fetal wellbeing, unless birth imminent

Management



Ref.236965

*Discuss with NICU

Date Issued: April 2017

Review Date: May 2019

Written/Authorised by: Maternity Guidelines Group

Review Team: Maternity Guidelines Group

Diagnosis and Management of Cervical Insufficiency GLM0055

Maternity Guidelines

Christchurch Women's Hospital

Christchurch New Zealand